

STIC-ILL

MIC
QH442.643

Fr m: Marx, Irene
Sent: Wednesday, July 31, 2002 8:53 AM
T : STIC-ILL
Subject: 09/842637
Imp rtance: High

Please send to Irene Marx, Art Unit 1651; CM1, Room 10E05, phone 308-2922, Mail box in 11B01

Courvalin et al., ASM News 1992, 58:368-375

Evers et al., 1993, Gene, 124:143-144

Quinn et al., 1989, Antimicrob. Agents Chemother. 33:1451-1456.

Rice et al., 1991, Antimicrob. Agents Chemother. 35:1243-1244

Sahm et al., 1989, J. Clin, Microbiol., 33:1588-1591

Sahm et al., 1990, Antimicrob. Agents Chemother. 34:1846.-1848

Neu et al., 1992, Science, 257:1064-1073

Jorgensen, 1993, Diagn. Microbiol. Infect Dis 16:245-249

Jorgensen et al., 1993, J. Clin, Microbiol., 31:2841-2844

Irene Marx
Art Unit 1651
CMI 10-E-05,
Mail Box 11-B-01
703-308-2922

GENE 06921

The *vanB* gene of vancomycin-resistant *Enterococcus faecalis* V583 is structurally related to genes encoding D-Ala:D-Ala ligases and glycopeptide-resistance proteins VanA and VanC*

(D-alanine:D-alanine ligase; cell wall; peptidoglycan synthesis)

Stefan Evers^a, Daniel F. Sahm^b and Patrice Courvalin^a

^aUnité des Agents Antibactériens, Institut Pasteur, 75724 Paris Cedex 15, France; and ^bJewish Hospital at Washington University Medical Center, Saint-Louis, MO 63110, USA

Received by F. Barany: 14 October 1992; Accepted: 19 October 1992; Received at publishers: 2 November 1992

SUMMARY

We report the cloning and sequencing of a 632-bp amplified fragment internal to the *vanB* gene of vancomycin-resistant (Vm^R) *Enterococcus* (*En.*) *faecalis* V583. The DNA fragment hybridized to Vm^R strains of *En. faecium* and *En. faecalis*, but not to their susceptible derivatives.

Glycopeptide antibiotics vancomycin (Vm) and teicoplanin (Te) bind to the C-terminal D-Ala residues of peptidoglycan precursors blocking their incorporation into the bacterial cell wall (Reynolds, 1989). These residues are incorporated into cell wall precursors as a dipeptide synthesized by D-Ala:D-Ala ligases (Ddl) (Walsh, 1989). The VanA ligase synthesizes the depsipeptide D-Ala-D-Lac which substitutes for D-Ala-D-Ala leading to synthesis of precursors which bind Vm with reduced

affinity (Bugg et al., 1991; Handwerger et al., 1992; Messer and Reynolds, 1992).

Glycopeptide resistance in enterococci is heterogeneous (Dutka-Malen et al., 1990). Resistance proteins

Correspondence to: Dr. P. Courvalin, Unité des Agents Antibactériens, 28, rue du Dr. Roux, 75724 Paris, Cedex 15, France. Tel. (33-1) 45 68 83 20; Fax (33-1) 45 68 83 19.

* On request, the authors will supply experimental evidence for the conclusions reached in this brief note.

Abbreviations: aa, amino acid(s); bp, base pair(s); D-Ala, D-alanine(s); DdlA and DdlB, D-Ala:D-Ala ligases of *E. coli*; D-Lac, D-lactate; *E.*, *Escherichia*; *En.*, *Enterococcus*; kb, kilobase(s) or 1000 bp; nt, nucleotide(s); oligo, oligodeoxyribonucleotide; PCR, polymerase chain reaction; ^R, resistant; ^S, sensitive; Te, teicoplanin; VanA, *En. faecium* Vm-resistance-conferring protein; VanB, *En. faecalis* Vm-resistance-conferring protein; VanC, *En. gallinarum* Vm-resistance-conferring protein; *ranB*, gene encoding VanB; Vm, vancomycin.

```
L F E L S G I P Y V G C D I O S S A A C . 20
TCTGTTGAATTGCTGGTATCCCTATGTAGGTCGGATATTCAAAGCTCCGAGCTTG 60
M D K S L A Y I L T K N A G I A V P E F 40
CATGGACAAATCACTGGCCTACATCTTACAAATAATCGGGCATCGCCGTCGCCGAAT 120
Q M I E R G D K P E A R T L T Y P V F V 60
TCAAATGATTGAAAAGGTGACAAACCGGAGCGAGGACGCTTACCTACCTGTCTTTGT 180
K P A R S G S S F G V T K V N S T E E L 80
GAAGCCGGCAGCGTCAGGTTCTGTCCTTTGGCGTAACCAAGTAACAGTACGGAAGAACT 240
N A A I E A A G O Y D G K I L I E Q A I 100
AAACGCTGCGATAGAGCAGCAGGACAATATGATGGAAAAATCTTAATTGAGCAAGCGAT 300
S G C E V G C A V M G N E D D L I V G E 120
TTGGGCTGTGAGGTCGGCTGCGCGGTCATGGGAACGAGGATGATTGATTGTCTCGGCA 360
V D G C I T L S H C I F R I H O E N E P E 140
AGTGGATCAATCCGGTTGAGCCAGGATATCTTCGCGATCCATCAGGAAAACGAGCGGA 420
K G S E N A M I V P A D I P V E E R N 160
AAAAGGCTCAGAGATGCGATGATTATCGTTCCAGCAGACATCCGGTCGAGGAACGAA 480
R V Q E T A K K V Y R V L G C R G L A R 180
TCGGGTGCAAGAAACGGCAAGAAAGTATATCGGGTCTTGGATGCAAGAGGCTTGCCTCG 540
V D L F L Q E D G G I V L E E V 196
TGTGATCTTTTTTTCAGGAGGATGCGGCATCGTTTAAACGAGGTC 589
```

Fig. 1. Nucleotide and corresponding aa sequence of the PCR fragment internal to the *ranB* gene. The nt sequence of both strands was determined from a pUC18 insert by the dideoxy-chain-termination method (Sanger et al., 1977) using T7 DNA polymerase. The sequences complementary to oligos V1 and V2 (Dutka-Malen et al., 1992) are not shown. Additional experiments were carried out to eliminate the possibility of nt misincorporation by the *Taq* DNA polymerase. GenBank accession No. is L06138.

VanB	LFELSGIPYV	GGDIQSSAAC	MDKSLAYILT	ENAGIAVPEF	QHIEKGRP	-----EA	RTITYPVFK	PARSGSSFGV	TKVNSTEELN	AAIEAAGQYD	GKILIEQAIS	101
VanA	LFELSGIPFV	GGDIQSSAIC	MDKSLTYIVA	KNAGIATPAF	WVINKDRP	-----VA	ATFTYPVFK	PARSGSSFGV	KKVNSEDELQ	YAIESAROYD	SKILIEQAVS	101
VanC	LLELMPLPV	GCHVAASALC	MXKMLHQLA	DTMGIASAPT	LLLSRYE	---PATIDRFI	QDHGFFPIK	PNEAGSSKGI	TKVDTKALQ	SALTAFAYG	STVLIQKAIA	107
DdlA	HLRVNLPFV	GSDVLSAAC	MDKDVTKRL	RDAGLNIAF	ITLTRANRHN	ISFAE---VE	SKLGLPLFVR	PARQSSVGV	SKVTSEEGYA	TAVALAFED	HKVIVEQGIN	107
DdlB	MLELNGLPYT	GSQVMASALS	MDKLRSKLLW	QGAGLPVAPH	VALTRAEFEK	GLSDKQLAEI	SALGLPVIVK	PSREGSSVGM	SKVVAENALQ	DALRLAFQHD	EEVLIEKMLS	110
	CC	CI	IC	C	II	CI	I	IC	II	IC	I	CCC
VanB	GCEVGCAMVG	NEDDLIVGEV	DQIRLSHGIF	RIHQENEPEK	GSENAIIVP	ADIPVEERNR	VQETAKKVYR	VLGCRGLARV	DLFLQEDGGI	VLNEV	196	
VanA	GCEVGCALVG	NSAALVGEV	DQIRLQYGIF	RIHQEVEPEK	GSENAVITVP	ADLSAEERGR	IQETAKKIYK	ALGCRGLARV	DMFLQDNGRI	VLNEV	196	
VanC	GIEIGCGILG	NE-QLTIGAC	DAISLVGFF	DFEERYQLIS	---ATITVP	APLPLALESQ	IKEQAQLLYR	NLCITGLARI	DFEVTNQGAI	YLNEI	197	
DdlA	GREIECAVLG	NDNP---QA	STCGEIVLTS	DFYAYDTKYI	DEGDAVVVP	AAIAPEINDK	IRAIIVQAYQ	TLCGAGMARV	DVFLTPEREV	VINEI	198	
DdlB	GPFTVAILG	EEIL-----	PSRIQPSG	TFYDYKAYL	SDETQYFC-P	AGLEASQAN	LQALVLKAWT	TLGCKGWGRI	DVMDSDGQF	YLLEA	198	
	I	IC	CCCCI		I	IC	C	II	I	CIC	ICCC	C

Fig. 2. Alignment of the deduced partial aa sequence of VanB and of the corresponding regions of VanA, VanC, DdlA and DdlB (Dutka-Malen et al., 1992). Identical aa (I) and conservative substitutions (C) in the five sequences are indicated below the alignment. For classification as conservative substitutions, the aa were grouped as follows: RK, LFPMVI, STQNC, AGW, H, ED and Y.

VanA and VanC display 28 to 38% aa identity with Ddl of *E. coli* (Dutka-Malen et al., 1992). The structural genes for VanA and VanC do not hybridize with DNA of enterococci that become resistant to Vm only after induction (VanB phenotype) (Dutka-Malen et al., 1990; Leclercq et al., 1992).

Oligos V1 and V2 allow PCR amplification of fragments internal to genes encoding VanA, VanC, and Ddl (Dutka-Malen et al., 1992). These oligos prime the amplification of ca. 600-bp fragments from *En. faecalis* V583 and *En. faecium* D366 which display the VanB phenotype (Sahm et al., 1989; Gutmann et al., 1992). The fragments from strain V583 were cloned into pUC18 (Norrander et al., 1983) and the insert of a recombinant plasmid was sequenced (Fig. 1). The deduced aa sequence of the insert was similar to a portion of VanA (77% aa identity), of VanC (37%) and of Ddl of *E. coli* (30 and 32%) (Fig. 2). In Southern hybridization, the cloned fragment hybridized with a 3.3-kb *HindIII-KpnI* fragment of *En. faecalis* V583 and a 7.5-kb *HindIII-KpnI* fragment of *En. faecium* D366 (data not shown). The probe did not hybridize to DNA from either Vm^s derivatives of these strains or Vm^s *En. faecalis* and *En. faecium* reference strains. These results suggest that the cloned PCR product corresponds to an internal fragment of a resistance-conferring gene acquired by the Vm^R strains. This gene encoded a Ddl-related enzyme, designated VanB, which could be involved in the synthesis of a substitute for D-Ala-D-Ala. This hypothesis is consistent with preliminary characterization of peptidoglycan precursors from *En. faecium* D366 (Billot-Klein et al., 1992).

ACKNOWLEDGEMENTS

We thank M. Arthur and S. Dutka-Malen for helpful discussions and for critical reading of the manuscript and L. Gutmann for the gift of bacterial strains.

REFERENCES

- Billot-Klein, D., Gutmann, L., Collatz, E. and Van Heijenoort, J.: Analysis of peptidoglycan precursors in vancomycin-resistant enterococci. *Antimicrob. Agents Chemother.* 36 (1992) 1487-1490.
- Bugg, T.D.H., Wright, G., Dutka-Malen, S., Arthur, M., Courvalin, P. and Walsh, C.T.: Molecular basis for vancomycin resistance in *Enterococcus faecium* BM4147: biosynthesis of a depsipeptide, peptidoglycan precursor by vancomycin resistant proteins VanH and VanA. *Biochemistry* 30 (1991) 10408-10415.
- Dutka-Malen, S., Leclercq, R., Coutant, V., Duval, J. and Courvalin, P.: Phenotypic and genotypic heterogeneity of glycopeptide resistance determinants in gram-positive bacteria. *Antimicrob. Agents Chemother.* 34 (1990) 1875-1879.
- Dutka-Malen, S., Molinas, C., Arthur, M. and Courvalin, P.: Sequence of the *vanC* gene of *Enterococcus gallinarum* BM4174 encoding a D-alanine:D-alanine ligase-related protein necessary for vancomycin resistance. *Gene* 112 (1992) 53-58.
- Gutmann, L., Billot-Klein, D., Al-Obeid, S., Klarc, I., Francoual, S., Collatz, E. and Van Heijenoort, J.: Inducible carboxypeptidase activity in vancomycin-resistant enterococci. *Antimicrob. Agents Chemother.* 36 (1992) 77-80.
- Handwerger, S., Pucci, M.J., Volk, K.J., Liu, J. and Lee, J.: The cytoplasmic peptidoglycan precursor of vancomycin-resistant *Enterococcus faecalis* terminates in lactate. *J. Bacteriol.* 174 (1992) 5982-5984.
- Leclercq, R., Dutka-Malen, S., Duval, J. and Courvalin, P.: Vancomycin resistance gene *vanC* is specific to *Enterococcus gallinarum*. *Antimicrob. Agents Chemother.* 36 (1992) 2005-2008.
- Messer, J. and Reynolds, P.: Modified peptidoglycan precursors produced by glycopeptide-resistant enterococci. *FEMS Microbiol. Lett.* 94 (1992) 195-200.
- Norrander, J., Kempe, T. and Messing, J.: Construction of improved M13 vectors using oligodeoxynucleotide-directed mutagenesis. *Gene* 26 (1983) 101-106.
- Reynolds, P.E.: Structure, biochemistry and mechanism of action of glycopeptide antibiotics. *Eur. J. Clin. Microbiol. Infect. Dis.* 8 (1989) 943-950.
- Sahm, D.F., Kissinger, J., Gilmore, M.S., Murray, P.R., Mulder, R., Solliday, J. and Clarke, B.: In vitro susceptibility studies of vancomycin-resistant *Enterococcus faecalis*. *Antimicrob. Agents Chemother.* 33 (1989) 1588-1591.
- Sanger, F., Nicklen, S. and Coulson, A.R.: DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74 (1977) 5463-5467.
- Walsh, C.T.: Enzymes in the D-alanine branch of bacterial cell-wall peptidoglycan assembly. *J. Biol. Chem.* 264 (1989) 2393-2396.